

## GMP APC anti-human CD79b (Ig $\beta$ ) Antibody

<b>Catalog# / Size</b>	260294 / 100 tests
<b>Clone</b>	CB3-1
<b>Workshop</b>	VI CD79.1
<b>Other Names</b>	Igb, Ig $\beta$ (Ig-beta), B29
<b>Isotype</b>	Mouse IgG1, $\kappa$
<b>Description</b>	CD79 is a heterodimeric molecule comprised of an $\alpha$ -chain (CD79a) and $\beta$ -chain (CD79b). A 37-39 kD type I integral membrane protein CD79b is non-covalently associated with CD79a and cell surface IgM to form the B-cell receptor (BCR) complex. CD79b is expressed on the surface of surface Ig (slg)-positive B cells and in the cytoplasm of slg-negative B cells. It is essential for signal transduction after surface Ig crosslinking.

### Product Details

---

<b>Reactivity</b>	Human
<b>Reported Reactivity</b>	African Green, Baboon, Cynomolgus, Rhesus
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA) and a stabilizer.
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
<b>Concentration</b>	100.0 $\mu$ g/mL
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 $\mu$ L per million cells in 100 $\mu$ L staining volume or 5 $\mu$ L per 100 $\mu$ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
<b>Excitation Laser</b>	Red Laser (633 nm)
<b>Application References</b>	1. Nakamura T, <i>et al.</i> 1992. <i>P. Natl. Acad. Sci. USA</i> 89:8522. 2. Garcia Vela J, <i>et al.</i> 1999. <i>Leukemia</i> . 13:1501. 3. Yoshino N, <i>et al.</i> 2000. <i>Exp. Anim. (Tokyo)</i> 49:97. (FC) 4. Rossi EA, <i>et al.</i> 2014. <i>PLoS One</i> . 9:98315. <a href="#">PubMed</a>
<b>(PubMed link indicates BioLegend citation)</b>	

<b>Disclaimer</b>	<p><b>GMP RUO Flow Cytometry Antibodies.</b> BioLegend GMP RUO fluorophore conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or therapeutic procedures. Our processes include:</p> <ul style="list-style-type: none"><li>• Batch-to-batch consistency</li><li>• Material traceability</li><li>• Documented procedures</li><li>• Documented employee training</li><li>• Equipment maintenance and monitoring records</li><li>• Lot-specific certificates of analysis</li><li>• Quality audits per ISO 13485:2016</li><li>• QA review of released products</li></ul>
-------------------	---

## Antigen Details

<b>Structure</b>	37-39 kD type I integral membrane protein, associate with CD79a and cell surface IgM to form the B-cell receptor (BCR) complex, Ig superfamily.
<b>Distribution</b>	B cells
<b>Function</b>	From BCR complex, signal transduction
<b>Cell Type</b>	B cells
<b>Biology Area</b>	Immunology
<b>Molecular Family</b>	CD Molecules
<b>Antigen References</b>	1. Zola H, et al. Eds. 2007. <i>Leukocyte and Stromal Cell Molecules: The CD Markers</i> . A John Wiley & Sons Inc. 2. Dylke J, et al. 2007. <i>Immunol. Lett.</i> 112:47
<b>Gene ID</b>	<a href="#">974</a>


## Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats

PE anti-human CD79b (Ig $\beta$ ), APC anti-human CD79b (Ig $\beta$ ), PerCP/Cyanine5.5 anti-human CD79b (Ig $\beta$ ), Alexa Fluor® 647 anti-human CD79b (Ig $\beta$ ), PE/Cyanine7 anti-human CD79b (Ig $\beta$ ), APC/Fire™ 750 anti-human CD79b (Ig $\beta$ ), TotalSeq™-A0187 anti-human CD79b (Ig $\beta$ ), TotalSeq™-C0187 anti-human CD79b (Ig $\beta$ ), TotalSeq™-B0187 anti-human CD79b (Ig $\beta$ ), TotalSeq™-D0187 anti-human CD79b (Ig $\beta$ ), PE anti-human CD79b, APC anti-human CD79b, PerCP/Cyanine5.5 anti-human CD79b

## Product Data



Typical results from human peripheral blood lymphocytes stained either with CB3-1 APC used at 5  $\mu$ L/test (filled histogram) or with an isotype control (open histogram).

For research use only. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses.

8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587