

GMP APC anti-human CD41 Antibody

Catalog# / Size	260358 / 100 tests
Clone	HIP8
Workshop	IV P38
Other Names	gpIIb, CD41a
Isotype	Mouse IgG1, κ
Description	CD41 is a 125/25 kD α subunit of the gpIIb/IIIa (CD41/CD61) complex. CD41 is a heterodimer composed of a heavy chain (gpIIb α) and light chain (gpIIb β) linked by a single disulfide bond. It is a member of the integrin family primarily expressed on platelets and megakaryocytes. CD41 has been reported to be involved with platelet aggregation and platelet attachment to the ECM. CD41/CD61 complex acts as the receptor for fibrinogen, fibronectin, Von Willebrand factor, and thrombin.

Product Details

Reactivity	Human
Reported Reactivity	African Green, Baboon, Capuchin Monkey, Cynomolgus, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA) and a stabilizer
Preparation	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
Concentration	25 μ g/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Red Laser (633 nm)
Application Notes	Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections and blocking of platelet aggregation ² . The HIP8 antibody has been reported to block the activation of platelets by various stimuli, including collagen, and ADP.
Application References	<ol style="list-style-type: none"> Knapp W, <i>et al.</i> 1989. Leucocyte Typing IV. Oxford University Press. New York. McCarty OJT, <i>et al.</i> 2000. <i>Blood</i> 96:1789. Yoshino N, <i>et al.</i> 2000. <i>Exp. Anim. (Tokyo)</i> 49:97. (FC) Zhi L <i>et al.</i> 2013. <i>PLoS One.</i> 8:e79869. (IHC)
(PubMed link indicates BioLegend citation)	
Disclaimer	<p>GMP RUO Flow Cytometry Antibodies. BioLegend GMP RUO fluorophore conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or therapeutic procedures. Our processes include:</p> <ul style="list-style-type: none"> • Batch-to-batch consistency • Material traceability • Documented procedures • Documented employee training • Equipment maintenance and monitoring records

- Lot-specific certificates of analysis
- Quality audits per ISO 13485:2016
- QA review of released products

Antigen Details

Structure	Integrin family, α subunit of CD41/CD61 (GPIIb-IIIa) complex, 125/22 kD
Distribution	Platelets, megakaryocytes
Function	Platelet aggregation, platelet attachment to extracellular matrix
Ligand/Receptor	Fibrinogen, fibronectin, von Willebrand factor, thrombin
Cell Type	Megakaryocytes, Platelets
Biology Area	Cell Adhesion, Cell Biology, Immunology
Molecular Family	Adhesion Molecules, CD Molecules
Antigen References	1. Denzin L, <i>et al.</i> 1996. <i>J. Exp. Med.</i> 184:2153. 2. Denzin L, <i>et al.</i> 1995. <i>Cell</i> 82:155. 3. Riberdy J, <i>et al.</i> 1994. <i>J. Cell Biol.</i> 125:1225.
Gene ID	3674


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-human CD41, FITC anti-human CD41, PE anti-human CD41, PE/Cyanine5 anti-human CD41, Purified anti-human CD41, Pacific Blue™ anti-human CD41, APC/Cyanine7 anti-human CD41, PE/Cyanine7 anti-human CD41, PerCP/Cyanine5.5 anti-human CD41, Purified anti-human CD41 (Maxpar® Ready), Alexa Fluor® 647 anti-human CD41, Alexa Fluor® 700 anti-human CD41, Alexa Fluor® 488 anti-human CD41, Brilliant Violet 421™ anti-human CD41, PE/Dazzle™ 594 anti-human CD41, Brilliant Violet 510™ anti-human CD41, Biotin anti-human CD41, TotalSeq™-A0353 anti-human CD41, TotalSeq™-C0353 anti-human CD41, Brilliant Violet 785™ anti-human CD41, Brilliant Violet 605™ anti-human CD41, Ultra-LEAF™ Purified anti-human CD41, TotalSeq™-B0353 anti-human CD41 Antibody, APC/Fire™ 750 anti-human CD41 Antibody, TotalSeq™-D0353 anti-human CD41

Product Data



Typical results from Human peripheral blood platelets stained with either HIP8 APC used at 5 μ L/test (filled histogram) or with an isotype control (open histogram).

For research use only. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all

applicable Limited Use Label Licenses.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com

Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587