

GMP APC anti-human IFN- γ Antibody

Catalog# / Size	260372 / 100 tests
Clone	4S.B3
Other Names	Interferon- γ , Immune interferon, Type II interferon, T cell interferon, Macrophage-activating factor (MAF), IFN-g, IFN-gamma
Isotype	Mouse IgG1, κ
Description	Interferon- γ is a potent multifunctional cytokine which is secreted primarily by activated NK cells and T cells. Originally characterized based on anti-viral activities, IFN- γ also exerts anti-proliferative, immunoregulatory, and proinflammatory activities. IFN- γ can upregulate MHC class I and II antigen expression by antigen-presenting cells.

Product Details

Reactivity	Human
Reported Reactivity	Chimpanzee, Baboon, Cynomolgus, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Partially purified, native human IFN- γ
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA) and a stabilizer.
Preparation	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
Concentration	25 μ g/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Red Laser (633 nm)
Application Notes	<p>ELISA or ELISPOT Detection⁵: The biotinylated 4S.B3 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with purified NIB42 antibody (Cat. No. 502402/502404) or purified MD-1 antibody (Cat. No. 507502/507513) as the capture antibody.</p> <p>Flow Cytometry^{3,4,6-8}: The fluorochrome-labeled 4S.B3 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IFN-γ-producing cells within mixed cell populations.</p> <p>Additional reported applications (for the relevant formats) include: neutralization^{1,2}, Western blotting, immunohistochemical staining of paraformaldehyde-fixed, saponin-treated tissue sections, and immunocytochemistry. The 4S.B3 antibody can neutralize the bioactivity of natural or recombinant IFN-γ.</p> <p>Note: For testing human IFN-γ in serum or plasma, BioLegend's ELISA Max™ Sets (Cat. No. 430101 to 430106) are specially developed and recommended.</p>

Application References

(PubMed link indicates BioLegend citation)

1. Meager A, *et al.* 1984. *J. Interferon Res.* 4:619. (Neut)
2. Meager A, 1987. *Lymphokines and Interferons: A Practical Approach.* IRL Press Ltd, Oxford, p. 105. (Neut)
3. Sester M, *et al.* 2002. *J. Virol.* 76:3748. (ICFC)
4. Infante-Duarte C, *et al.* 2000 *J. Immunol.* 165:6107. (ICFC)
5. Goodier M, *et al.* 2000. *J. Immunol.* 165:139. (ELISA)
6. Chen H, *et al.* 2005. *J. Immunol.* 175:591. (ICFC)
7. Smeltz RB, 2007. *J. Immunol.* 178:4786. (ICFC)
8. Iwamoto S, *et al.* 2007. *J. Immunol.* 179:1449. (ICFC) [PubMed](#)
9. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (ICFC)

Disclaimer

GMP RUO Flow Cytometry Antibodies. BioLegend GMP RUO fluorophore conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or therapeutic procedures. Our processes include:

- Batch-to-batch consistency
- Material traceability
- Documented procedures
- Documented employee training
- Equipment maintenance and monitoring records
- Lot-specific certificates of analysis
- Quality audits per ISO 13485:2016
- QA review of released products

Antigen Details

Structure	Cytokine; dimer; 20-25 kD (Mammalian)
Bioactivity	Antiviral/antiparasitic activities; inhibits proliferation; enhances MHC class I and II expression on APC
Cell Sources	CD8 ⁺ and CD4 ⁺ T cells, NK cells
Cell Targets	T cells, B cells, macrophages, NK cells, endothelial cells, fibroblasts
Receptors	IFN- γ R α (CDw119) dimerized with IFN- γ R β (AF-1)
Cell Type	Tregs
Biology Area	Cell Biology, Immunology, Neuroinflammation, Neuroscience
Molecular Family	Cytokines/Chemokines
Antigen References	<ol style="list-style-type: none">1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook.</i> Academic Press, San Diego.2. De Maeyer E, <i>et al.</i> 1992. <i>Curr. Opin. Immunol.</i> 4:321.3. Farrar M, <i>et al.</i> 1993. <i>Annu. Rev. Immunol.</i> 11:571.4. Gray P, <i>et al.</i> 1987. <i>Lymphokines</i> 13:151.
Regulation	Upregulated by IL-2, FGF-basic, EGF; downregulated by vitamin D3 or DMN; labile at pH2
Gene ID	3458

Related Protocols


- [Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

PE anti-human IFN- γ , APC anti-human IFN- γ , FITC anti-human IFN- γ , Biotin anti-human IFN- γ , Purified anti-human IFN- γ , Alexa Fluor® 488 anti-human IFN- γ , Alexa Fluor® 647 anti-human IFN- γ , Alexa Fluor® 700 anti-human IFN- γ , Pacific Blue™ anti-human IFN- γ , PerCP/Cyanine5.5 anti-human IFN- γ , APC/Cyanine7 anti-human IFN- γ , PE/Cyanine7 anti-human IFN- γ , Brilliant Violet 421™ anti-human IFN- γ , Brilliant Violet 570™ anti-human IFN- γ , Brilliant Violet 605™ anti-human IFN- γ , Brilliant Violet 650™ anti-human IFN- γ , Brilliant Violet 711™ anti-human IFN- γ , Brilliant Violet 785™ anti-human IFN- γ , Brilliant Violet 510™ anti-human IFN- γ , PE/Dazzle™ 594 anti-human IFN- γ , APC/Fire™ 750 anti-human IFN- γ , PerCP anti-human IFN- γ , Brilliant Violet 750™ anti-

human IFN- γ , KIRAVIA Blue 520™ anti-human IFN- γ Antibody, Spark NIR™ 685 anti-human IFN- γ Antibody, Spark Blue™ 515 anti-human IFN- γ , Spark UV™ 387 anti-human IFN- γ , Spark PLUS UV™ 395 anti-human IFN- γ

Product Data



Typical result of PMA + Ionomycin with Brefeldin A stimulated (4-hour) human peripheral blood lymphocytes fixed with Fixation Buffer (Cat# 420801), permeabilized with Permeabilization Wash Buffer (Cat# 421002), and intracellularly stained either with 4S.B3 APC used at 5 μ L/test (filled histogram) or with an isotype control (open histogram).

For Research Use Only. Suitable for *ex vivo* cell processing. Not for injection or diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587