

GMP PE/Dazzle™ 594 anti-human CD7 Antibody

Catalog# / Size	260380 / 100 tests
Clone	CD7-6B7
Workshop	IV T-164
Other Names	gp40
Isotype	Mouse IgG2a, κ
Description	CD7 is a 40 kD type I transmembrane glycoprotein also known as gp40. It is a member of the immunoglobulin superfamily found on T cells, NK cells, thymocytes, hematopoietic progenitors, and monocytes (weakly). CD7 is also expressed on acute lymphocytic leukemia (ALL) and some acute myeloid leukemia (AML) cells. CD7 crosslinking induces a calcium flux in T lymphocytes, presumably as a result of cytoplasmic domain association with PI3-kinase. CD7 costimulation can induce cytokine secretion and modulate cellular adhesion.

Product Details

Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	KG1a cell line
Formulation	Phosphate-buffered solution, pH 7.2, containing True-Stain Monocyte Blocker™, 0.09% sodium azide and 0.2% (w/v) BSA (origin USA) and a stabilizer.
Preparation	The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions.
Concentration	50 µg/mL
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. * PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	Additional reported (for the relevant formats) applications include proteogenomics ² .
Application References (PubMed link indicates BioLegend citation)	<ol style="list-style-type: none">1. Knapp W, <i>et al.</i> 1989. <i>Leucocyte Typing IV: White Cell Differentiation Antigens</i>. Oxford University Press.2. Peterson VM, <i>et al.</i> 2017. <i>Nat. Biotechnol.</i> 35:936. (PG)

Disclaimer

GMP RUO Flow Cytometry Antibodies. BioLegend GMP RUO fluorophore conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or therapeutic procedures. Our processes include:

- Batch-to-batch consistency
- Material traceability
- Documented procedures
- Documented employee training
- Equipment maintenance and monitoring records
- Lot-specific certificates of analysis
- Quality audits per ISO 13485:2016
- QA review of released products

Antigen Details

Structure	Ig superfamily, type I transmembrane glycoprotein, 40 kD
Distribution	T cells, NK cells, thymocytes, hematopoietic progenitors, myeloid leukemic cell subsets
Function	T cell activation
Ligand/Receptor	Cytoplasmic domain associates with P13-kinase
Cell Type	Hematopoietic stem and progenitors, Leukemia, NK cells, T cells, Thymocytes
Biology Area	Costimulatory Molecules, Immunology
Molecular Family	CD Molecules
Antigen References	1. Barclay N, <i>et al.</i> 1993. <i>The Leucocyte Antigen FactsBook</i> . Academic Press Inc. San Diego. 2. Stillwell R, <i>et al.</i> 2001. <i>Immunol. Res.</i> 24:31. 3. Rabinowich H, <i>et al.</i> 1994. <i>J. Immunol.</i> 152:517.
Gene ID	924


Related Protocols

- [Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human CD7, FITC anti-human CD7, PE anti-human CD7, APC anti-human CD7, PE/Cyanine5 anti-human CD7, PE/Cyanine7 anti-human CD7, Purified anti-human CD7 (Maxpar® Ready), PerCP/Cyanine5.5 anti-human CD7, PE/Dazzle™ 594 anti-human CD7, APC/Fire™ 750 anti-human CD7, Alexa Fluor® 647 anti-human CD7, TotalSeq™-A0066 anti-human CD7, Alexa Fluor® 700 anti-human CD7, TotalSeq™-C0066 anti-human CD7, TotalSeq™-B0066 anti-human CD7, Pacific Blue™ anti-human CD7 Antibody, TotalSeq™-D0066 anti-human CD7, GMP APC anti-human CD7, GMP FITC anti-human CD7, Pacific Blue™ anti-human CD7, GMP PE anti-human CD7, GMP PE/Cyanine7 anti-human CD7, GMP Pacific Blue™ anti-human CD7, Spark NIR™ 685 anti-human CD7

Product Data



Typical results from human peripheral blood lymphocytes stained either with CD7-6B7 PE/Dazzle™ 594 used at 5 μ L/test (filled histogram) or with an isotype control (open histogram).

For Research Use Only. Suitable for *ex vivo* cell processing. Not for injection or diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587