

## GMP APC anti-human CD79a (Igα) Antibody

<b>Catalog# / Size</b>	260392 / 100 tests
<b>Clone</b>	HM47
<b>Workshop</b>	V cB017
<b>Other Names</b>	Mb-1, Iga, CD79
<b>Isotype</b>	Mouse IgG1, κ
<b>Description</b>	CD79a is a 47 kD type I integral membrane protein, also known as mb-1 or Iga. It is a member of the Ig superfamily and disulphide-associated with CD79b (B29). The interaction of CD79a/CD79b heterodimer with B cell surface Ig forms B cell antigen receptor complex. CD79a is expressed in B cells from early pre-B to plasma cell stage. It has been shown that CD79a is also weakly expressed in some precursors of T- and myeloid cells. CD79 mediates the transport of IgM to B cell surface and transduces signals initiated by BCR aggregation.

### Product Details

---

<b>Reactivity</b>	Human
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA) and a stabilizer
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.
<b>Concentration</b>	25 µg/mL
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">ICFC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">intracellular immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
<b>Excitation Laser</b>	Red Laser (633 nm)
<b>Disclaimer</b>	<b>GMP RUO Flow Cytometry Antibodies.</b> BioLegend GMP RUO fluorophore conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or therapeutic procedures. Our processes include: <ul style="list-style-type: none"><li>• Batch-to-batch consistency</li><li>• Material traceability</li><li>• Documented procedures</li><li>• Documented employee training</li><li>• Equipment maintenance and monitoring records</li><li>• Lot-specific certificates of analysis</li><li>• Quality audits per ISO 13485:2016</li><li>• QA review of released products</li></ul>

### Antigen Details

---

<b>Structure</b>	47 kD, type I integral protein, Ig superfamily, CD79a/CD79b heterodimer interacts with B cell membrane Ig.
<b>Distribution</b>	B cells (from early pre-B to plasma cell stage), some T and myeloid cell precursors
<b>Function</b>	In association with CD79b and B cell membrane Ig to form B cell antigen receptor complex
<b>Cell Type</b>	B cells, Hematopoietic stem and progenitors
<b>Biology Area</b>	Immunology
<b>Molecular Family</b>	CD Molecules
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Zola Heddy, <i>et al.</i> Eds. 2007. <i>Leukocyte and Stromal Cell Molecules: The CD markers</i>. WILEY-LISS</li> <li>2. Astsaturov IA, <i>et al.</i> 1996. <i>Leukemia</i> 10:769</li> <li>3. Mson DY, <i>et al.</i> 1995 <i>Blood</i> 86:1453</li> <li>4. Hashimoto M, <i>et al.</i> 2002. <i>J. Pathol.</i> 197:341</li> </ol>
<b>Gene ID</b>	<a href="#">973</a>


## Related Protocols

- [Intracellular Flow Cytometry Staining Protocol](#)

## Other Formats

Purified anti-human CD79a (Ig $\alpha$ ), PE anti-human CD79a (Ig $\alpha$ ), APC anti-human CD79a (Ig $\alpha$ ), PerCP/Cyanine5.5 anti-human CD79a (Ig $\alpha$ ), PE/Cyanine7 anti-human CD79a (Ig $\alpha$ ), FITC anti-human CD79a (Ig $\alpha$ ), Alexa Fluor® 594 anti-human CD79a (Ig $\alpha$ ), Alexa Fluor® 647 anti-human CD79a (Ig $\alpha$ ), APC anti-human CD79a, PerCP/Cyanine5.5 anti-human CD79a, PE anti-human CD79a, FITC anti-human CD79a

## Product Data



Typical results from human peripheral blood lymphocytes were fixed with Fixation Buffer (Cat# 420801), permeabilized with Permeabilization Wash Buffer (Cat# 421002), and intracellularly stained with HM47 APC used at 5  $\mu$ L/test (filled histogram) or with an isotype control (open histogram).

For Research Use Only. Suitable for *ex vivo* cell processing. Not for injection or diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at [www.biolegend.com/terms](http://www.biolegend.com/terms) ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587