

GMP PE anti-human IL-4 Antibody

Catalog# / Size	260400 / 100 tests
Clone	MP4-25D2
Other Names	Interleukin-4, Ia inducing factor (IaIF), B-cell stimulating factor-1 (BSF-1), Hodgkin's cell growth factor (HCGF), Mast cell growth factor-2 (MCGF-2), Macrophage fusion factor (MFF), T cell growth factor-2 (TCGF-2)
Isotype	Rat IgG1, κ
Description	IL-4 is a pleiotropic cytokine that is produced by activated T cells, mast cells, and basophils. IL-4 elicits many different biological responses but has two dominant functions. The first is regulating differentiation of naïve CD4 ⁺ T cell to the Th2 type. Th2 cells produce IL-4, IL-5, IL-10, and IL-13, which tend to favor a humoral immune response while suppressing a cell-mediated immune response controlled by Th1 cells. The second is regulating IgE and IgG1 production by B cells.

Product Details

Reactivity	Human
Reported Reactivity	Pig, Rhesus
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	CHO-expressed, recombinant human IL-4
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA) and a stabilizer.
Preparation	The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions.
Concentration	25 $\mu\text{g}/\text{mL}$
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μL per million cells in 100 μL staining volume or 5 μL per 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
Application Notes	ELISA Detection ^{1,3} or ELISPOT Detection ^{4,5} : The biotinylated MP4-25D2 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with purified 8D4-8 antibody (Cat. No. 500702/500707) as the capture antibody. Flow Cytometry ^{6,9} : The fluorochrome-labeled MP4-25D2 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IL-4 -producing cells within mixed cell populations. Neutralization ¹⁻³ : The LEAF™ purified antibody (Endotoxin <0.1 EU/ μg , Azide-Free, 0.2 μm filtered) is recommended for neutralization of human IL-4 bioactivity (Cat. No. 500815). The MP4-25D2 antibody can neutralize the bioactivity of natural or recombinant IL-4.

Application References

(PubMed link indicates BioLegend citation)

1. Chretien I, *et al.* 1989. *J. Immunol. Methods* 117:67. (ELISA Detection, Neut)
2. Ramanathan L, *et al.* 1993. *Biochem.* 32:3549. (Neut)
3. Abrams J, *et al.* 1992. *Immunol. Rev.* 127:5. (ELISA Detection, Neut)
4. Mahanty S, *et al.* 1992. *J. Immunol.* 148:3567. (ELISPOT Detection)
5. Klinman D, *et al.* 1994. *Curr. Prot. Immunol.* John Wiley and Sons New York. Unit 6.19. (ELISPOT Detection)
6. Prussin C, *et al.* 1995. *J. Immunol. Methods* 188:117. (ICFC)
7. Raqib R, *et al.* 1995. *Infect. Immun.* 63:289.
8. Andersson J, *et al.* 1994. *Immunology* 83:16.
9. Iwamoto S, *et al.* 2007. *J. Immunol.* 179:1449. (ICFC) [PubMed](#)
10. Kubota M, *et al.* 1997. *J. Immunol.* 158:5321.
11. Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113. [PubMed](#)
12. Kroneke MA, *et al.* 2012. *J. Immunol.* 188:3734. [PubMed](#)

Disclaimer

GMP RUO Flow Cytometry Antibodies. BioLegend GMP RUO fluorophore conjugated antibodies are manufactured in a dedicated GMP facility and compliant with ISO 13485:2016. For research use only. Not for use in diagnostic or therapeutic procedures. Our processes include:

- Batch-to-batch consistency
- Material traceability
- Documented procedures
- Documented employee training
- Equipment maintenance and monitoring records
- Lot-specific certificates of analysis
- Quality audits per ISO 13485:2016
- QA review of released products

Antigen Details

Structure	Cytokine; 15-19 kD (Mammalian)
Bioactivity	Differentiation of naïve CD4 ⁺ T cells to the T _H 2 type, proliferation/differentiation of activated B cells, expression of class II MHC antigens, and of low affinity IgE receptors in resting B cells
Cell Sources	Mast cells, T cells, bone marrow stromal cells
Cell Targets	B cells, T cells, monocytes, endothelial cells, fibroblasts
Receptors	Heterodimer IL-4R α (CD124); γ -subunit (CD132) in common with IL-2R, IL-7R, IL-13R, IL-15R
Cell Type	Tregs
Biology Area	Cell Biology, Immunology, Neuroinflammation, Neuroscience
Molecular Family	Cytokines/Chemokines
Antigen References	<ol style="list-style-type: none">1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook</i>. Academic Press San Diego.2. Boulay J, <i>et al.</i> 1992. <i>Curr. Opin. Immunol.</i> 4:294.3. Dullens H, <i>et al.</i> 1991. <i>In vivo</i> 5:567.4. Paul W. 1991. <i>Blood</i> 77:1859.
Regulation	Upregulated by IL-2, platelet activating factor; downregulated by TGF- β
Gene ID	3565

Related Protocols


- [Intracellular Flow Cytometry Staining Protocol](#)

Other Formats

APC anti-human IL-4, Biotin anti-human IL-4, FITC anti-human IL-4, PE anti-human IL-4, Purified anti-human IL-4, Alexa Fluor® 488 anti-human IL-4, Alexa Fluor® 647 anti-human IL-4, Brilliant Violet 421™ anti-human IL-4, PerCP/Cyanine5.5 anti-human IL-4, PE/Cyanine7 anti-human IL-4, Brilliant Violet 605™ anti-human IL-4, Purified anti-human IL-4 (Maxpar® Ready), PE/Dazzle™ 594

anti-human IL-4, APC/Cyanine7 anti-human IL-4, Brilliant Violet 510™ anti-human IL-4, Ultra-LEAF™ Purified anti-human IL-4, TotalSeq™-B1149 anti-human IL-4, Brilliant Violet 785™ anti-human IL-4, Brilliant Violet 711™ anti-human IL-4

Product Data



Typical result of PMA + Ionomycin with Brefeldin A stimulated (4-hour) human peripheral blood lymphocytes fixed with Fixation Buffer (Cat# 420801), permeabilized with Permeabilization Wash Buffer (Cat# 421002), and intracellularly stained either with MP4-25D2 PE used at 5 μ L/test (left) or with an isotype control (right).

For Research Use Only. Suitable for *ex vivo* cell processing. Not for injection or diagnostic or therapeutic use.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587