

PerCP/Cyanine5.5 anti-human CD33

Analyte Specific Reagent. Analytical and performance characteristics are not established.

Catalog# / Size 983906 / 500 μL

Clone WM53

Workshop IV M-505

Other Names Siglec-3, gp67, p67

Isotype Mouse IgG1, κ

Description CD33 is a 67 kD type I transmembrane glycoprotein also known as Siglec-3, gp67, and p67. It

is a sialoadhesion immunoglobulin superfamily member expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells. CD33 is absent on normal platelets, lymphocytes, erythrocytes and hematopoietic stem cells. CD33 functions as a sialic acid-

dependent cell adhesion molecule with carbohydrate/lectin binding activity.

Product Details

Reactivity Human

Formulation Phosphate-buffered solution, pH 7.2, containing True-Stain Monocyte Blocker™, 0.09% sodium

azide and 0.2% (w/v) BSA (origin USA).

Preparation The antibody was purified by affinity chromatography, and conjugated with PerCP/Cyanine5.5

under optimal conditions

Concentration 200 µg/mL

Storage & Handling The antibody solution should be stored undiluted between 2°C and 8°C and protected from


prolonged exposure to light. Do not freeze.

Application Suggested for Flow Cytometry

Disclaimer WARNINGS AND PRECAUTIONS

- Use appropriate personal protective equipment and safety practices per universal precautions when working with this reagent. Refer to the reagent safety data sheet.
- This antibody contains sodium azide. Follow federal, state and local regulations to dispose of this reagent. Sodium azide build-up in metal wastepipes may lead to explosive conditions; if disposing of reagent down wastepipes, flush with water after disposal.
- 3. All specimens, samples and any material coming in contact with them should be considered potentially infectious and should be disposed of with proper precautions and in accordance with federal, state and local regulations.
- 4. Do not use this reagent beyond the expiration date stated on the label.
- 5. Do not use this reagent if it appears cloudy or if there is any change in the appearance of the reagent as these may be indication of possible deterioration.
- 6. Avoid prolonged exposure of the reagent or stained cells to light.

Product Data



Typical results from human peripheral blood monocytes stained either with WM53 PerCP/Cyanine5.5 used at 5 μL/test (filled histogram) or with an isotype control (open histogram).

Symbols Glossary*

Symbols clossery							
Symbol	Meaning	Symbol Title	Symbol No.	Symbol	Meaning	Symbol Title	Symbol No.
REF	Catalog number	Catalogue number	5.1.6	(i	Indicates the need for the user to consult the instructions for use.	Consult instructions for use	5.4.3
1	Indicates the temperature limits to which the medical device can be safely exposed.	Temperature limit	5.3.7	*	Indicates a medical device that needs protection from light sources.	Keep away from sunlight	5.3.2
K	Indicates the upper limit of temperature to which the medical device can be safely exposed.	temperature	5.3.6	Ω	Indicates the date after which the medical device is not to be used.	Use-by date	5.1.4
•••	Indicates the medical device manufacturer.	Manufacturer	5.1.1	EC REP	Indicates the authorized representative in the European Community.	Authorized representative in the European Community	
	Indicates the manufacturer's	Batch code	5.1.5		Indicates a medical device that is	In vitro diagnostic	5.5.1
LOT	batch code so that the batch or lot can be identified.			IVD	intended to be used as an in vitro diagnostic medical device.	medical device	

^{*} Symbol information is from EN ISO 15223-1:2016 Medical devices – Symbols to be used with medical device labels, labelling and information to be supplied - Part 1: General requirements

Analyte Specific Reagent. Analytical and performance characteristics are not established.

This product is supplied subject to the terms and conditions, including the limited license, located at www.biolegend.com/terms ("Terms") and may be used only as provided in the Terms. Without limiting the foregoing, BioLegend products may not be used for any Commercial Purpose as defined in the Terms, resold in any form, used in manufacturing, or reverse engineered, sequenced, or otherwise studied or used to learn its design or composition without express written approval of BioLegend. Regardless of the information given in this document, user is solely responsible for determining any license requirements necessary for user's intended use and assumes all risk and liability arising from use of the product. BioLegend is not responsible for patent infringement or any other risks or liabilities whatsoever resulting from the use of its products.

BioLegend, the BioLegend logo, and all other trademarks are property of BioLegend, Inc. or their respective owners, and all rights are reserved.

8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587