



**Purified anti-mouse CD105 Antibody**



C57BL/6 mouse frozen spleen section was fixed with 4% paraformaldehyde (PFA) for ten minutes at room temperature and blocked with 5% FBS for 30 minutes at room temperature. Then the section was stained with 10 µg/ml of purified anti-mouse CD105 (clone MJ7/18) and 10 µg/ml of Alexa Fluor<sup>®</sup> 647 anti-mouse CD3ε (clone 145-2C11) (green) overnight at 4°C, followed by 2.5 µg/ml of Alexa Fluor<sup>®</sup> 594 Goat anti-rat IgG (clone Poly4054) (red) for two hours at room temperature. Nuclei were counterstained with DAPI (blue). The image was captured with a 10X objective.

## **Purified anti-mouse CD105 Antibody**



C57BL/6 mouse frozen liver section was fixed with 4% paraformaldehyde (PFA) for ten minutes at room temperature and blocked with 5% FBS for 30 minutes at room temperature. Then the section was stained with 10  $\mu\text{g}/\text{ml}$  of purified anti-mouse CD105 (clone MJ7/18) overnight at 4°C, followed by 2.5  $\mu\text{g}/\text{ml}$  of Alexa Fluor<sup>®</sup> 594 Goat anti-rat IgG (clone Poly4054) (red) for two hours at room temperature. Nuclei were counterstained with DAPI (blue). The image was captured with a 10X objective.